



## Rubisco

**Root growth enhancer, and regulator of foliar transpiration.**

**Prevents and cures drought and salinity.**

**N 11+ Fe-B-Cu-Zn-Mn-Mo**

### BENEFITS

- Rich in antioxidants and sea weeds high in cytokines and betaines. Increases, strengthens and deepens root mass. Relieves roots damages caused by maintenance errors
- Stimulates the rapid installation of pre-cultivated sod and, after laying, its abundant rooting
- Prevents all stresses and damages related to scarce water availability. Recovers turf suffering from drought. Prevents localized dry spot ; you save on wetting agents
- Regulates stomata opening for optimal foliar transpiration and xylem water transport. This allows proper plant's cooling. Reduces irrigation in the summer up to 30%, producing drier and firmer playing turf areas

### TECHNICAL FEATURES

11% Total nitrogen (N) 1% nitrogen from amino acids (L); 10% urea nitrogen  
0.01% Boron (B) soluble in water  
0.002 Copper (Cu) water soluble EDTA  
0.02% Iron (Fe) water soluble, EDTA  
0.01% Manganese (Mn) water soluble EDTA  
0.002% Zinc (Zn) water soluble, EDTA  
0.001% Molybdenum (Mo) water soluble  
Contains L-Amino Acids of vegetal origin (proline)  
Mannitol, seaweed meal filtrate. Contains: organic foliar absorption facilitators

Roots enhancer and foliar antistress fertilizer, manufactured with a proprietary formula derived from University of Bologna research on the activators of the enzymes involved in regulating water trans-cellular movements (osmo protectors) and, in particular, the chemical elements involved in the regulation of stomata opening and closing during evapo-transpiration. Rich in antioxidants and sea weeds with a very high cytokinin and betaine content, promotes roots growth in all periods of the year

### APPLICATION METHODS

0,5 liters/1.000 m<sup>2</sup> in 100 liters of water. A 5 liter drum is sufficient for 10.000 m<sup>2</sup> of surface. Distribute on dry grass, preferentially in mid morning when stomata are fully opened ( temperature between 15 and 25°C ). Do not irrigate or mow for 6 hours after treatment. Avoid contact of product on sidewalks or other stone surfaces

**PACKAGING:** 5 liter drums and 1 liter bottles



ON THE TURF'S SIDE